Brandon J. Barnes, M.D. | Orthopaedic Surgeon, Sports Medicine Specialist www.osswf.com (239) 334-7000

ACL Reconstruction with Quadriceps Tendon Autograft and Meniscal Repair (All Inside) Protocol

| | Weight-baring | Brace | Range of Motion | Exercises |
|-----------------------|--|---|-----------------------------|--|
| Phase 1 0-6 weeks | Full in brace (a) | 0-2 week: Locked in full extension for ambulation and sleeping | Passive 0-90° by 2 weeks | Calf pumps, patellar mobilization, quadriceps tightening sets and SLR in brace. No weight-bearing motion. |
| | | Off for exercises and hygiene only | | Add side-lying hip and core, advance quad set and stretching (a) |
| | | 2-6 weeks: Unlocked 0-90°for ambulation, remove for sleeping (a) | | In brace: Addition of heel raises, total gym (closed chain), gait normalization, eccentric quads, eccentric hamstrings |
| | | | | Advance core work with hip, gluteal and core strengthening |
| | | | | No weight-bearing with flexion >90° |
| Phase 2 6-12 weeks | Full, progressing to normal gait pattern | Discontinue at 6 weeks if quadriceps control is adequate (no extension lag) | Gain full and pain- free | Begin toe raises, closed chain quads, balance exercises, hamstring curls, stationary bike, step-ups, front and side planks, hip/glute/core |
| Phase 3 | Full with a | None | Full | Advance closed chain strengthening |
| 12-16 weeks | normalized gait pattern | | | Progress proprioception activities |
| | | | | Begin stair climber, elliptical at 8 weeks and running straight at 12 weeks when 8" step down is satisfactory |
| Phase 4 | Full | None | Full | 16 weeks: Begin jumping |
| 16 weeks – | | | | 20 weeks: Advance running to sprinting, backward running, cutting, pivoting, changing |

| 6 months | | | | direction, initiate plyometric program and sport- specific drills |
|-------------|------|------|------|--|
| | | | | 22 weeks: Advance as tolerated |
| | | | | FSA completed at 22 weeks (b) |
| Phase 5 | Full | None | Full | Gradual return to sports participation after |
| 6-12 months | | | | completion of functional sports assessment (FSA) |
| | | | | Maintenance program based on FSA |

SLR - Straight-leg raise

- (a) Modified with concomitantly performed meniscus repair/transplantation or articular cartilage procedure
- (b) Completion of FSA (Functional Sports Assessment) is not mandatory, but is recommended at approximately 22 weeks post- op for competitive athletes returning to play after rehabilitation