

Post-Op Instructions for De Quervain's Release

After your surgery, post-surgical discomfort or pain is likely. This discomfort can last several days to a few weeks. At certain times of the day your discomfort may be more intense. Start range of motion exercises including making a fist and touching your thumb to all your fingers as tolerated to prevent stiffness.

Dressings:

You will have a splint in place that will remain on until your first post operative visit. Keep the splint clean and dry. Do not soak the incision (no baths, pool) for 2-3 weeks.


For pain you have after your surgery we recommend the following:

1. If you were prescribed narcotic pain medication, take it for moderate to severe pain only.
2. Take **Ibuprofen** 800mg by mouth every 8 hours for 3 days after surgery, then take as needed. (Do not take Ibuprofen if you have been instructed by your primary doctor not to take NSAIDs or anti-inflammatories due to Kidney problems, history of stomach ulcers, or if you are also on a blood thinner or anti-coagulant like lovenox, coumadin, xarelto, etc.)
3. Take **Tylenol** (acetaminophen) regularly alternating with the Ibuprofen. There is typically acetaminophen in the narcotic pain medication, so be sure not to exceed 3000 mg of acetaminophen per 24 hours.
4. Focus on **elevation**, keeping the hand at the level of the heart or higher. When up and walking around, you should have the hand resting on your chest. When sitting, you can rest the arm on the elbow with fingers pointing toward the ceiling or rest the hand on several pillows.
5. **Nausea is a common side effect of many pain medications. You will want to eat something before taking your pain medicine to help prevent nausea.**
6. If needed, using a **cold pack** to ice the affected area a few times a day (15 to 20 minutes at a time) can help to relieve pain, reduce swelling and bruising.

3 weeks post operatively


Now that you are 3 weeks post operatively, you should have most if not all of your range of motion back, but you will need to continue doing your tendon gliding exercises for at least the next 3 weeks, as your body continues to form scar.

Tendon Gliding Exercises



Perform this sequence of fists 10 repetitions, 5 times per day.

Now that your incision is healed and we can be more vigorous with your wrist motion, it is important to glide the median nerve along its path to avoid it also being surrounded by scar tissue. That is done by performing this series of motions:



Repeat this sequence 5 repetitions, 5 times per day – Hold each position 10 seconds.

Post operative instructions:

Scar Care

Massage your scar with lotion for 5 minutes 5 times per day. Use a moderately deep pressure to help flatten the scar and decrease its sensitivity to pressure.

Returning to Activities

Most patients can resume full use of their hand after surgery by about 6 weeks post operatively, but please let your comfort be your guide. Do not repetitively perform activities that cause you pain. A gradual return to activities without pain is our goal.

EMERGENCIES - Contact Dr. Barnes' office at (239) 334-7000 if any of the following are present:

- Painful swelling or numbness (note that some swelling and numbness is normal)
- Unrelenting pain
- Fever (over 101° F – it is normal to have a low grade fever (<100°) for the first day or two following surgery) or chills.
- Redness around incisions
- Continuous drainage or bleeding from incision (a small amount of drainage is expected)
- Difficulty breathing
- Excessive nausea/vomiting

If you have an emergency that requires immediate attention proceed to the nearest emergency room.