## Brandon J. Barnes, M.D. | Orthopaedic Surgeon, Sports Medicine Specialist www.osswf.com

(239) 334-7000

## Distal Femoral Osteotomy (DFO) Protocol

	Weight-baring	Brace	Range of Motion	Exercises
Phase 1 0-2 weeks	Heel touch only	Locked in full extension at all times	Gentle passive 0- 90°	Calf pumps, patellar mobilization, quadriceps tightening sets and SLR in brace
		Off for hygiene and home exercise only		
Phase 2	Heel touch only	2-8 weeks: Unlocked 0-	Maintain full	Progress non-weight-bearing flexibility,
2-6 weeks		until 6 weeks progress to full	extension and	modalities as needed
			flexion	Add side-lying hip and core, advance quad set and stretching
		Off at night for sleeping	IIEXION	
		Discontinue brace at 6 weeks		Advance core work with hip, gluteal and core strengthening
Phase 3	Advance 25% weekly and progress to full with normalized gait pattern	None	Full	Progress closed chain activities
6-8 weeks				Advance hamstring exercises, proprioception & balance exercises, lunges & leg press
				Advance core work with hip, gluteal and core strengthening
				Begin Stationary Bike at 6 weeks
Phase 4	Full	None	Full	Progress Phase III exercises and
8-16 weeks				<b>functional activities</b> : walking lunges, planks, bridges, Swiss ball, half-Bosu exercises
				Advance core work with hip, gluteal and core strengthening

				Add elliptical at 12 weeks and swimming permitted at 12 weeks
Phase 5	Full	None	Full	Advance Phase 4 activities
4-6 months				Progress to functional training, including impact activity after 5 months when cleared by MD
CPM – contir	uous passive m	otion machine; SLR – Straig	ht-leg raise	