Brandon J. Barnes, M.D. | Orthopaedic Surgeon, Sports Medicine Specialist BrandonBarnesMD.com (239) 334-7000

Meniscus Root Repair Protocol

	Weight-baring	Brace	Range of Motion	Exercises
Phase 1 0-4 weeks	Non weight bearing with crutches	Locked in full extension at all times	Gentle passive 0-90°	Calf pumps, patellar mobilization, quadriceps tightening sets and SLR in brace. No weight-bearing motion.
		Off for hygiene and home exercise only		
		Off at night at 2 weeks		
Phase 2 4-8 weeks	4-6 weeks: 50% weight bearing with brace	4-6 weeks: Unlocked 0-90°	Advance motion to regain full passive motion	Add side-lying hip and core, advance quad set and stretching (a)
4-0 WEEKS	unlocked 0-90° 6-8 weeks: Progress to full with brace unlocked 0-90° and discontinue brace at 8 weeks.	Off at night Discontinue brace at 8 weeks or transition to unloader brace at 6 weeks		In brace: Addition of heel raises, total gym (closed chain), gait normalization, eccentric quads, eccentric hamstrings.
				Advance core work with hip, gluteal and core strengthening
				No weight-bearing with flexion >90°
Phase 3	Full	None	Full	Progress closed chain activities
8-12 weeks				Advance hamstring exercises, proprioception & balance exercises, lunges & leg press 0-90° only
				Advance core work with hip, gluteal and core strengthening
				Begin Stationary Bike at 8 weeks
Phase 4 3-5 months	Full	None	Full	Progress Phase III exercises and functional activities: walking lunges, planks, bridges, Swiss ball, half-Bosu exercises

		Advance core work with hip, gluteal and core strengthening Swimming allowed at 12 weeks		
		Advance to running/jumping and sport specific drills at 16 weeks when cleared by MD		
SLD Straight leg raise				

SLR – Straight-leg raise

(a) Avoid tibial rotation for first 6 weeks.