**Brandon J. Barnes, M.D.** | Orthopaedic Surgeon, Sports Medicine Specialist BrandonBarnesMD.com (239) 334-7000

## **Microfracture or BioCartilage Protocol**

	Weight-baring	Brace	Range of Motion	Exercises
Phase 1 0-6 weeks	Heel touch only	O-2 weeks: Locked in full extension at all times  Off for hygiene and home exercise only  Discontinue after 2 weeks	CPM 6 hrs/day; begin 0-40° and advance 5-10° daily as tolerated Gentle passive 0-90°	Calf pumps, patellar mobilization, quadriceps tightening sets and SLR in brace  2-6 weeks: Add side-lying hip and core, advance quadriceps and hamstring sets
Phase 2 6-8 weeks	Advance 25% weekly and progress to full with normalized gait pattern	None	Full	Advance core work with hip, gluteal and core strengthening
Phase 3 8-12 weeks	Full	None	Full	Gait normalization with closed chain activities: wall sits, shuttle, mini-squats, toe raises.  Advance core work with hip, gluteal and core strengthening  Begin unilateral stance activities and balance training
Phase 4 3-6 months	Full	None	Full	Progress Phase III exercises and functional activities: walking lunges, planks, bridges, Swiss ball, half-Bosu exercises  Advance core work with hip, gluteal and core strengthening  May advance to bike, elliptical, pool as tolerated

Phase 5	Full	None	Full	Advance non-impact activity			
6-12 Months				No running, jumping, pivoting, or sports participation unless cleared by MD (after 8 months)			
CPM – continuous passive motion machine; SLR – Straight-leg raise							